

## ASAM Criteria Clinical Decision-Making for the New Professional (and a Refresher for Others)


David Mee-Lee, M.D. Davis, CA  
(530) 753-4300; Voice mail: (916) 715-5856  
davidmlee@gmail.com www.changecompanies.net  
www.trainforchange.net www.tipsttopics.com  
www.ASAMCriteria.org www.instituteforwellness.com

Breakout Session April 18, 2017 12:45 PM-1:45 PM Des Moines, IA  
40<sup>th</sup> Annual Governor's Conference on Substance Abuse

### A. Underlying Principles of the ASAM Criteria - Generations of Clinical Care


#### (a) Complications-driven Treatment

- ⤴ No diagnosis of Substance Use Disorder
- ⤴ Treatment of complications of addiction with no continuing care
- ⤴ Relapse triggers treatment of complications only




#### (b) Diagnosis, Program-driven Treatment


- ⤴ Diagnosis determines treatment
- ⤴ Treatment is the primary program and aftercare
- ⤴ Relapse triggers a repeat of the program



#### (c) Individualized, Clinically-driven Treatment



d) Client-Directed, Outcome-Informed Treatment (Feedback Informed Treatment)



2. Assessment of Biopsychosocial Severity and Function (The ASAM Criteria 2013, pp 43-53)

The common language of six ASAM Criteria dimensions determine needs/strengths:

1. Acute intoxication and/or withdrawal potential
2. Biomedical conditions and complications
3. Emotional/behavioral/cognitive conditions and complications
4. Readiness to Change
5. Relapse/Continued Use/Continued Problem potential
6. Recovery environment

Assessment Dimensions	Assessment and Treatment Planning Focus
1. Acute Intoxication and/or Withdrawal Potential	Assessment for intoxication and/or withdrawal management. Withdrawal management in a variety of levels of care and preparation for continued addiction services
2. Biomedical Conditions and Complications	Assess and treat co-occurring physical health conditions or complications. Treatment provided within the level of care or through coordination of physical health services
3. Emotional, Behavioral or Cognitive Conditions and Complications	Assess and treat co-occurring diagnostic or sub-diagnostic mental health conditions or complications. Treatment provided within the level of care or through coordination of mental health services
4. Readiness to Change	Assess stage of readiness to change. If not ready to commit to full recovery, engage into treatment using motivational enhancement strategies. If ready for recovery, consolidate and expand action for change
5. Relapse, Continued Use or Continued Problem Potential	Assess readiness for relapse prevention services and teach where appropriate. If still at early stages of change, focus on raising consciousness of consequences of continued use or problems with motivational strategies.
6. Recovery Environment	Assess need for specific individualized family or significant other, housing, financial, vocational, educational, legal, transportation, childcare services

### 3. Biopsychosocial Treatment - Overview: 5 M's

- \* Motivate - Dimension 4 issues; engagement and alliance building
- \* Manage - the family, significant others, work/school, legal
- \* Medication – withdrawal management; HIV/AIDS; MAT - anti-craving anti-addiction meds; disulfiram, methadone; buprenorphine, naltrexone, acamprosate, psychotropic medication
- \* Meetings - AA, NA, Al-Anon; SMART Recovery, Dual Recovery Anonymous, etc.
- \* Monitor - continuity of care; relapse prevention; family and significant others

### 4. Treatment Levels of Service (*The ASAM Criteria* 2013, pp 106-107)

- 1 Outpatient Services
- 2 Intensive Outpatient/Partial Hospitalization Services
- 3 Residential/Inpatient Services
- 4 Medically-Managed Intensive Inpatient Services

<b>ASAM Criteria Level of Withdrawal Management Services for Adults</b>	<b>Level</b>	<b>Note: There are no separate Withdrawal Management Services for Adolescents</b>
Ambulatory Withdrawal Management without Extended On-Site Monitoring	1-WM	Mild withdrawal with daily or less than daily outpatient supervision; likely to complete withdrawal management and to continue treatment or recovery
Ambulatory Withdrawal Management with Extended On-Site Monitoring	2-WM	Moderate withdrawal with all day WM support and supervision; at night, has supportive family or living situation; likely to complete WM.
Clinically-Managed Residential Withdrawal Management	3.2-WM	Moderate withdrawal, but needs 24-hour support to complete WM and increase likelihood of continuing treatment or recovery
Medically-Monitored Inpatient Withdrawal Management	3.7-WM	Severe withdrawal and needs 24-hour nursing care and physician visits as necessary; unlikely to complete WM without medical, nursing monitoring
Medically-Managed Inpatient Withdrawal Management	4-WM	Severe, unstable withdrawal and needs 24-hour nursing care and daily physician visits to modify WM regimen and manage medical instability
<b>ASAM Criteria Levels of Care</b>	<b>Level</b>	<b>Same Levels of Care for Adolescents except Level 3.3</b>
Early Intervention	0.5	Assessment and education for at risk individuals who do not meet diagnostic criteria for Substance-Related Disorder
Outpatient Services	1	Less than 9 hours of service/week (adults); less than 6 hours/week (adolescents) for recovery or motivational enhancement therapies/ strategies
Intensive Outpatient	2.1	9 or more hours of service/week (adults); 6 or more hours/week (adolescents) to treat multidimensional instability
Partial Hospitalization	2.5	20 or more hours of service/week for multidimensional instability not requiring 24 hour care
Clinically-Managed Low-Intensity Residential	3.1	24 hour structure with available trained personnel; at least 5 hours of clinical service/week
Clinically Managed Population-Specific High-Intensity Residential Services (Adult criteria only)	3.3	24 hour care with trained counselors to stabilize multidimensional imminent danger. Less intense milieu and group treatment for those with cognitive or other impairments unable to use full active milieu or therapeutic community
Clinically-Managed High-Intensity Residential	3.5	24 hour care with trained counselors to stabilize multidimensional imminent danger and prepare for outpatient treatment. Able to tolerate and use full active milieu or therapeutic community
Medically-Monitored Intensive Inpatient	3.7	24 hour nursing care with physician availability for significant problems in Dimensions 1, 2 or 3. Sixteen hour/day counselor ability
Medically-Managed Intensive Inpatient	4	24 hour nursing care and daily physician care for severe, unstable problems in Dimensions 1, 2 or 3. Counseling available to engage patient in treatment
Opioid Treatment Services	OTS	Opioid Treatment Program (OTP) – agonist meds: methadone, buprenorphine; Office Based Opioid Treatment (OBOT); antagonist medication - naltrexone

## D. Understanding Dimension 4, Readiness to Change

### 1. Developing the Treatment Contract *(The ASAM Criteria 2013, page 58)*

	<u>Client</u>	<u>Clinical Assessment</u>	<u>Treatment Plan</u>
<u>What?</u>	What does client want?	What does client need?	What is the Tx contract?
<u>Why?</u>	Why now? What's the level of commitment?	Why? What reasons are revealed by the assessment data?	Is it linked to what client wants?
<u>How?</u>	How will s/he get there?	How will you get him/her to accept the plan?	Does client buy into the link?
<u>Where?</u>	Where will s/he do this?	Where is the appropriate setting for treatment? What is indicated by the placement criteria?	Referral to level of care
<u>When?</u>	When will this happen? How quickly? How badly does s/he want it?	When? How soon? What are realistic expectations? What are milestones in the process?	What is the degree of urgency? What is the process? What are the expectations of the referral?

### 2. Stages of Change and How People Change -Transtheoretical Model (Prochaska & DiClemente)

**Pre-contemplation:** not yet considering the possibility of change although others are aware of a problem; active resistance to change; seldom appear for treatment without coercion; could benefit from non-threatening information to raise awareness of a possible “problem” and possibilities for change.

**Contemplation:** ambivalent, undecided, vacillating between whether he/she really has a “problem” or needs to change; wants to change, but this desire exists simultaneously with resistance to it; may seek professional advice to get an objective assessment; motivational strategies useful at this stage, but aggressive or premature confrontation provokes strong resistance and defensive behaviors; many Contemplators have indefinite plans to take action in the next six months or so.

**Preparation:** takes person from decisions made in Contemplation stage to the specific steps to be taken to solve the problem in the Action stage; increasing confidence in the decision to change; certain tasks that make up the first steps on the road to Action; most people planning to take action within the very next month; making final adjustments before they begin to change their behavior.

**Action:** specific actions intended to bring about change; overt modification of behavior and surroundings; most busy stage of change requiring the greatest commitment of time and energy; care not to equate action with actual change; support and encouragement still very important to prevent drop out and regression in readiness to change.


**Maintenance:** sustain the changes accomplished by previous action and prevent relapse; requires different set of skills than were needed to initiate change; consolidation of gains attained; not a static stage and lasts as little as six months or up to a lifetime; learn alternative coping and problem-solving strategies; replace problem behaviors with new, healthy life-style; work through triggers of relapse.

**Relapse and Recycling:** expectable, but not inevitable setbacks; avoid becoming stuck, discouraged, or demoralized; learn from relapse before committing to a new cycle of action; comprehensive, multidimensional assessment to explore all reasons for relapse.

**Termination:** this stage is the ultimate goal for all changers; person exits the cycle of change, without fear of relapse; debate over whether certain problems can be terminated or merely kept in remission through maintenance strategies.

**E. Imminent Danger** (*The ASAM Criteria* 2013, pp. 65-58) - Three components:

1. A strong probability that certain behaviors (such as continued alcohol or other drug use or addictive behavior relapse) will occur.
2. The likelihood that such behaviors will present a significant risk of serious adverse consequences to the individual and/or others (as in reckless driving while intoxicated, or neglect of a child).
3. The likelihood that such adverse events will occur in the very near future, within hours and days, rather than weeks or months



(*The ASAM Criteria* 2013, p 124)

---

## Tracy

A 16-year-old young woman is brought into the emergency room of an acute care hospital. She had gotten into an argument with her parents and ended up throwing a chair. There was some indication that she was intoxicated at the time and her parents have been concerned about her coming home late and mixing with the wrong crowd. There has been a lot of family discord and there is mutual anger and frustration between the teen and especially her father. No previous psychiatric or addiction treatment.

The parents are both present at the ER, but the police who had been called by her mother brought her. The ER physician and nurse from the psychiatric unit who came from the unit to evaluate the teen, both feel she needs to be in hospital given the animosity at home, the violent behavior and the question of intoxication. Using the six ASAM assessment dimensions, the biopsychosocial clinical data is organized as follows:

Dimension 1, Intoxication/Withdrawal: though intoxicated at home not long before the chair-throwing incident, she is no longer intoxicated and has not been using alcohol or other drugs in large enough quantities for long enough to suggest any withdrawal danger.

Dimension 2, Biomedical Conditions/Complications: she is not on any medications, has been healthy physically and has no current complaints

Dimension 3, Emotional/Behavioral/Cognitive: complex problems with the anger, frustration and family discord; chair throwing incident this evening, but is not impulsive at present in the ER.

Dimension 4, Readiness to Change: willing to talk to therapist; blames her parents for being overbearing and not trusting her; agrees to treatment, but doesn't want to be at home at least for tonight.

Dimension 5, Relapse/Continued Use/Continued Problem Potential: high likelihood that if released to go back home immediately, there would be a reoccurrence of the fighting and possibly violence again, at least with father.

Dimension 6, Recovery Environment: parents frustrated and angry too; mistrustful of patient; and want her in the hospital to cut down on the family fighting

Severity Profile:                      Dimension:      1        2        3        4        5        6  
Severity:

Services Needed:

Site of Care:

### 2. Assessing Severity and Level of Function (*The ASAM Criteria* 2013, pp 54-56)

To determine the multidimensional severity or level of function profile, consider each of the six ASAM ASAM Criteria dimensions as regards pertinent assessment data organized under the three H's - History, Here and Now, How Worried Now.

The *History* of a client's past signs, symptoms and treatment is important, but never overrides the *Here and Now* of how a client is presenting currently in signs and symptoms. e.g., if a person has by History had severe alcohol withdrawal with seizures, but has not been drinking Here and Now at a rate or quantity that would predict any significant withdrawal; and as you look at them, they are not shaky or in withdrawal so you are not Worried about severe withdrawal - then there is no significant Dimension 1 severity.

The *Here and Now* presentation of a client's current information of substance use and mental health signs and symptoms can override the *History* e.g., if a person has never had serious suicidal behavior before by History; and in the Here and Now is indeed depressed and impulsively suicidal, you would not dismiss their severe suicidality just because they had never done anything serious before. Especially if you talked with them now and you are *Worried* that they could not reach out to someone if they became impulsive, then the Dimension 3 severity would be quite high.

*How Worried Now* you are as the clinician, counselor or assessor determines your severity or level of function (LOF) rating for each ASAM dimension. The combination of the three H's: History; Here and Now; and How Worried Now guides the clinician in presenting the severity and LOF profile.

**Ann**

DSM-5 Diagnosis: Alcohol Use Disorder, severe; and Cannabis Use Disorder, moderate; Major Depression

Ann, a 32-year-old white, divorced female, came in for assessment for the first time ever. She has been abstinent for 48 hours from alcohol and reports that she has remained so far up to 72 hours during the past three months. When she has done this she states she has experienced sweats, internal tremors and nausea, but has never hallucinated, experienced D.T.'s or seizures.

She states she is in good health except for alcoholic hepatitis for which she was just released from the hospital one week ago. Her doctor referred her for assessment. She smokes up to 3 or 4 joints a day, but stopped yesterday. In addition to the above, Ann describes two past suicide attempts using sleeping pills, but the most recent attempt was three years ago and she sees a psychiatrist once a month for review of her medication. She takes Prozac for the depression and doesn't report abuse of her medication.

Ann reported that she lives in a rented apartment and has very few friends since moving away after her divorce a year ago. She is currently unemployed after being laid off when the supermarket she worked at closed. She has worked as a waitress, check-out person and sales person before and says she has never lost a job due to addiction.

Ann appears slightly anxious, but is not flushed. She speaks calmly and is cooperative. Ann shows awareness of her consequences from chemical use, but tends to minimize it and blame others including her ex-husband who left her without warning. She doesn't know much about alcoholism/chemical dependency, but wants to learn more. She has one son, age 11, who doesn't see any problems with her drinking and doesn't know about her marijuana use.

**F. Gathering Data on Policy and Payment Barriers** (*The ASAM Criteria* 2013, p 126)

- ▲ Policy, payment and systems issues cannot change quickly. However, as a first step towards reframing frustrating situations into systems change, each incident of inefficient or in adequate meeting of a client's needs can be a data point that sets the foundation for strategic planning and change
- ▲ Finding efficient ways to gather data as it happens in daily care can provide hope and direction for change

**PLACEMENT SUMMARY**

<b>Level of Care/Service Indicated</b> - Insert the ASAM Level number that offers the most appropriate level of care/service that can provide the service intensity needed to address the client's current functioning/severity; and/or the service needed e.g., shelter, housing, vocational training, transportation, language interpreter	
<b>Level of Care/Service Received</b> - ASAM Level number -- If the most appropriate level or service is not utilized, insert the most appropriate placement or service available and circle the Reason for Difference between Indicated and Received Level or Service	
<b>Reason for Difference</b> - Circle only one number -- 1. Service not available; 2. Provider judgment; 3. Client preference; 4. Client is on waiting list for appropriate level; 5. Service available, but no payment source; 6. Geographic accessibility; 7. Family responsibility; 8. Language; 9. Not applicable; 10. Not listed (Specify):	
<b>Anticipated Outcome If Service Cannot Be Provided</b> – Circle only one number - 1. Admitted to acute care setting; 2. Discharged to street; 3. Continued stay in acute care facility; 4. Incarcerated; 5. Client will dropout until next crisis; 6. Not listed (Specify):	

---

## **LITERATURE REFERENCES**

“Addiction Treatment Matching – Research Foundations of the American Society of Addiction Medicine (ASAM) Criteria” Ed. David R. Gastfriend has released 2004 by The Haworth Medical Press. David Gastfriend edited this special edition that represents a significant body of work presented in eight papers. The papers address questions about nosology, methodology, and population differences and raise important issues to continually refine further work on the ASAM PPC. (To order: 1-800-HAWORTH; or [www.haworthpress.com](http://www.haworthpress.com))

Mee-Lee D, Shulman GD (2014): “The ASAM Criteria and Matching Patients to Treatment”, Chapter 27 in Section 4, Overview of Addiction Treatment in "Principles of Addiction Medicine" Eds Richard K. Ries, David A Fiellin, Shannon Miller, Richard Saitz. Fifth Edition. Lippincott Williams & Wilkins, Philadelphia, PA.,USA. pp 428 -441.

Mee-Lee D, Shulman GD, Fishman MJ, and Gastfriend DR, Miller MM eds. (2013). The ASAM Criteria: Treatment Criteria for Addictive, Substance-Related, and Co-Occurring Conditions. Third Edition. Carson City, NV: The Change Companies.

For more information on the new edition: [www.ASAMcriteria.org](http://www.ASAMcriteria.org)

Mee-Lee, David with Jennifer E. Harrison (2010): “Tips and Topics: Opening the Toolbox for Transforming Services and Systems”. The Change Companies, Carson City, NV

Mee-Lee, David (2013): “The ASAM Criteria: Getting Up to Date.” Advances in Addiction & Recovery Fall 2013. Vol.1., No. 3 pp 20-21.

Mee-Lee, David (2013): “The New Edition of the ASAM Criteria: What’s New and Why” Counselor Vol. 14. No.4 pp18-19. August 2013.

Mee-Lee, David (2013): “How to Really Use the New Edition of *The ASAM Criteria*: What To Do & What Not To Do.” Counselor Vol. 14. No.6 pp34-40. December 2013.

Prochaska, JO; Norcross, JC; DiClemente, CC (1994): “Changing For Good” Avon Books, New York.

## **RESOURCE FOR E-LEARNING AND INTERACTIVE JOURNALS**

E-learning module on “ASAM Multidimensional Assessment” and “From Assessment to Service Planning and Level of Care” – 5 CE credits for each module . “Introduction to The ASAM Criteria” (2 CEU hours) “Understanding the Dimensions of Change” – Creating an effective service plan” – Interactive Journaling “Moving Forward” – Guiding individualized service planning” – Interactive Journaling

To order: The Change Companies at 888-889-8866; [www.ASAMcriteria.org](http://www.ASAMcriteria.org)

## **CLIENT WORKBOOKS AND INTERACTIVE JOURNALS**

The Change Companies’ MEE (Motivational, Educational and Experiential) Journal System provides Interactive journaling for clients. It provides the structure of multiple, pertinent topics from which to choose; but allows for flexible personalized choices to help this particular client at this particular stage of his or her stage of readiness and interest in change.

To order: The Change Companies at 888-889-8866. [www.changecompanies.net](http://www.changecompanies.net).

## **FREE MONTHLY NEWSLETTER**

“TIPS and TOPICS” – Three sections: Savvy, Skills and Soul and at times additional sections: Stump the Shrink; Success Stories and Sharing Solutions. Sign up on [www.tipsntopics.com](http://www.tipsntopics.com) and click on “Subscribe”.